


# Improving quantitative precipitation forecasting by using high-resolution ensembles in PREVIEW



Universität Karlsruhe (TH)  
Forschungsuniversität • gegründet 1825

J.W. Schipper\*, M. Kunz, and Ch. Kottmeier



Institute for Meteorology and Climate Research/Forschungszentrum Karlsruhe, University of Karlsruhe, Germany

## Introduction

Effective operational flood management requires detailed knowledge of the distribution of precipitation at high spatial and temporal resolution. Within the interdisciplinary EU-IP PREVIEW (PREvention, Information, and Early Warning), it is aimed to achieve more precise flood forecasts through a better identification of precipitation fields and, in turn, to develop adequate and well-adapted prevention measures by working closely together with hydrologists.

## Goal

The goal is to provide information about the amount of precipitation during an extreme weather event. It is aimed to perform high resolution simulations (eventually at 1 km) resulting in a more realistically distributed precipitation because of a more realistic representation of orographic features. The simulations are performed with the standard weather forecasting model (LM) at 7 km resolution and the newly developed local model (LMK) at 2.8 and 1 km of the German Weather Service (DWD).

## General set-up

High-resolution model simulations will not exactly describe the atmosphere and, as a result, will not be able to predict the amount of precipitation in a certain area exactly. One approach to add some information about the stability to the simulations is to run several model simulation using slightly altered initial conditions each time. To avoid high computational costs, an ensemble of model simulations is run at a low resolution first, whereas the extreme precipitation relevant ones are rerun at high resolution (Fig. 1).




Figure 1: Graphical design of the iterative use of a set of ensembles to quantify the precipitation for extreme weather events.




Figure 2: Part of Europe with embedded research areas: data availability (a), low resolution – 7 km (b), high resolution simulations – 2.8 km (c).

## Research area

This paper focuses on the precipitation distribution in the German part of the river Danube catchment located in the south-eastern part of Germany covering roughly 450x450 km (Fig. 2 – panel c). The usual time scale of a flooding event in such a catchment is several hours to a few days, which acknowledges the urge for an effective operational flood-forecasting system. Within the framework of PREVIEW, close cooperation with hydrological groups is exhibited to gain better understanding of such a system.




Figure 3: Example of a simulated precipitation event (10/08/2002 11:00 local time) at a low (7 km) and a high (2.8 km) resolution.




Figure 4: The orography in the mountainous part of the research area at various resolutions (a: 7 km; b: 2.8 km; c: 1 km; d: 50 m)

## First results

First results show a significant difference in precipitation distribution between the simulations at a resolution of 7 km and 2.8 km (Fig. 3 – left and right panel, resp.). One of the reasons for this difference is the much finer resolved orography in the 2.8 km simulations (Fig. 4). It is expected that future simulations at a resolution of 1 km will give different results as well. These simulations will focus more on the river Iller in Bavaria (Germany) and will be able to better locate local maxima during extreme precipitation events.

## References

- Atger, F., 1999: The skill of ensemble prediction systems. *Mon. Wea. Rev.*, **127**, 1941–1953.
- Sattler, K., and Feddersen, H., 2005: Limited-area short-range ensemble predictions targeted for heavy rain in Europe. *Hydr. Earth Syst. Sci.*, **9**, 300–312

## Acknowledgements

This study is performed within the EU-IP PREVIEW. The authors thanks the German Weather Service (DWD) for giving helpful advice in case of the LMK simulations.